

A CLASS OF SIMPLE C^* -ALGEBRAS ARISING FROM CERTAIN NONSOFTIC SUBSHIFTS

KENGO MATSUMOTO

ABSTRACT. We present a class of subshifts $Z_N, N = 1, 2, \dots$ whose associated C^* -algebras $\mathcal{O}_{Z_N}$ are simple, purely infinite and not stably isomorphic to any Cuntz-Krieger algebra nor to Cuntz algebra. The class of the subshifts is the first examples whose associated C^* -algebras are not stably isomorphic to any Cuntz-Krieger algebra nor to Cuntz algebra. The subshifts Z_N are coded systems whose languages are context free. We compute the topological entropy for the subshifts and show that KMS-state for gauge action on the associated C^* -algebra $\mathcal{O}_{Z_N}$ exists if and only if the logarithm of the inverse temperature is the topological entropy for the subshift Z_N , and the corresponding KMS-state is unique.

Simple C^* -algebra, subshift, entropy, KMS-state.
Primary 46L35, 37B10; Secondary 54H20, 54C70

1. INTRODUCTION

Let Σ be a finite set with its discrete topology, that is called an alphabet. Each element of Σ is called a symbol. Let $\Sigma^{\mathbb{Z}}$ be the infinite product space $\prod_{i \in \mathbb{Z}} \Sigma_i$, where $\Sigma_i = \Sigma$, endowed with the product topology. The transformation σ on $\Sigma^{\mathbb{Z}}$ given by $\sigma((x_i)_{i \in \mathbb{Z}}) = (x_{i+1})_{i \in \mathbb{Z}}$ is called the full shift over Σ . Let Λ be a shift invariant closed subset of $\Sigma^{\mathbb{Z}}$ i.e. $\sigma(\Lambda) = \Lambda$. The topological dynamical system $(\Lambda, \sigma|_{\Lambda})$ is called a subshift or a symbolic dynamical system. It is written as Λ for brevity. Theory of symbolic dynamical system gives a basic method to study general dynamical systems (cf.[23]). It also has significant uses in coding for information theory. There is a class of subshifts called sofic shifts, that contains the topological Markov shifts. Sofic shifts are presented by finite square matrices with entries in formal sums of symbols. Such a matrix is called a symbolic matrix. It is an equivalent object to a finite labeled graph called a λ -graph. In [28], the author has introduced the notions of symbolic matrix system and λ -graph system. They are presentations of subshifts and generalizations of symbolic matrices and λ -graphs respectively. A symbolic matrix system $(\mathcal{M}, I)$ consists of a sequence of pairs $(\mathcal{M}_{l,l+1}, I_{l,l+1}), l \in \mathbb{Z}_+$ of rectangular symbolic matrices $\mathcal{M}_{l,l+1}$ and rectangular $\{0, 1\}$ -matrices $I_{l,l+1}$, where $\mathbb{Z}_+$ denotes the set of all nonnegative integers. Both the matrices $\mathcal{M}_{l,l+1}$ and $I_{l,l+1}$ have the same size for each $l \in \mathbb{Z}_+$. The column size of $\mathcal{M}_{l,l+1}$ is the same as the row size of $\mathcal{M}_{l+1,l+2}$. They satisfy the following commutation relations as symbolic matrices

$$I_{l,l+1} \mathcal{M}_{l+1,l+2} = \mathcal{M}_{l,l+1} I_{l+1,l+2}, \quad l \in \mathbb{Z}_+. \quad (1.1)$$

We further assume that for i there exists j such that the (i, j) -component $I_{l,l+1}(i, j) = 1$, and for j there uniquely exists i such that $I_{l,l+1}(i, j) = 1$. A λ -graph system $\mathfrak{L} = (V, E, \lambda, \iota)$ consists of a vertex set $V = V_0 \cup V_1 \cup V_2 \cup \dots$, an edge set

$E = E_{0,1} \cup E_{1,2} \cup E_{2,3} \cup \cdots$, a labeling map $\lambda : E \rightarrow \Sigma$ and a surjective map $\iota_{l,l+1} : V_{l+1} \rightarrow V_l$ for each $l \in \mathbb{Z}_+$. An edge $e \in E_{l,l+1}$ has its source vertex $s(e)$ in V_l , its terminal vertex $t(e)$ in V_{l+1} and its label $\lambda(e)$ in Σ . For a symbolic matrix system $(\mathcal{M}, I)$, the labeled edges from a vertex $v_i^l \in V_l$ to a vertex $v_j^{l+1} \in V_{l+1}$ are given by the symbols appearing in the (i, j) -component $\mathcal{M}_{l,l+1}(i, j)$ of $\mathcal{M}_{l,l+1}$. The matrix $I_{l,l+1}$ defines a surjection $\iota_{l,l+1}$ from V_{l+1} to V_l for each $l \in \mathbb{Z}_+$. By this observation, the symbolic matrix systems and the λ -graph systems are the same objects. They give rise to subshifts by taking the set of all label sequences appearing in the labeled Bratteli diagram. For a symbolic matrix system $(\mathcal{M}, I)$, let $M_{l,l+1}$ be the nonnegative rectangular matrix obtained from $\mathcal{M}_{l,l+1}$ by setting all the symbols equal to 1 for each $l \in \mathbb{Z}_+$. Then the resulting pair (M, I) satisfies the following relations by (1.1)

$$I_{l,l+1}M_{l+1,l+2} = M_{l,l+1}I_{l+1,l+2}, \quad l \in \mathbb{Z}_+. \quad (1.2)$$

We call (M, I) the nonnegative matrix system for $(\mathcal{M}, I)$. Let $m(l)$ be the row size of the matrix $I_{l,l+1}$ for each $l \in \mathbb{Z}_+$. Let $\mathbb{Z}_{I^t}$ be the abelian group defined by the inductive limit $\mathbb{Z}_{I^t} = \varinjlim \{I_{l,l+1}^t : \mathbb{Z}^{m(l)} \rightarrow \mathbb{Z}^{m(l+1)}\}$. The sequence $M_{l,l+1}^t, l \in \mathbb{Z}_+$ of the transposes of $M_{l,l+1}$ naturally acts on $\mathbb{Z}_{I^t}$ by the relation (1.2), that is denoted by $\lambda_{(M,I)}$. The K-groups for (M, I) have been defined in [28] as:

$$K_0(M, I) = \mathbb{Z}_{I^t} / (\text{id} - \lambda_{(M,I)})\mathbb{Z}_{I^t}, \quad K_1(M, I) = \text{Ker}(\text{id} - \lambda_{(M,I)}) \text{ in } \mathbb{Z}_{I^t}.$$

In [32], C^* -algebra $\mathcal{O}_{\mathfrak{L}}$ associated with a λ -graph system $\mathfrak{L}$ has been introduced. The C^* -algebras are generalizations of the Cuntz-Krieger algebras and the C^* -algebras associated with subshifts ([9], [24], cf. [4]). They are universal unique concrete C^* -algebras generated by finite families of partial isometries and sequences of projections subject to certain operator relations encoded by structure of the λ -graph systems. Let (M, I) be the nonnegative matrix system for the symbolic matrix system $(\mathcal{M}, I)$ of $\mathfrak{L}$. The K-theory formulae for $\mathcal{O}_{\mathfrak{L}}$ have been obtained as in the following way:

$$K_0(\mathcal{O}_{\mathfrak{L}}) = K_0(M, I), \quad K_1(\mathcal{O}_{\mathfrak{L}}) = K_1(M, I). \quad (1.3)$$

There is a canonical method to construct a λ -graph system $\mathfrak{L}$ from a given subshift Λ . The λ -graph system is called the canonical λ -graph system for Λ and written as $\mathfrak{L}^\Lambda$. The C^* -algebra $\mathcal{O}_{\mathfrak{L}^\Lambda}$ associated with the canonical λ -graph system $\mathfrak{L}^\Lambda$ coincides with the C^* -algebra $\mathcal{O}_\Lambda$ associated with subshift Λ (cf. [4], [24]). If in particular Λ is a topological Markov shift Λ_A for a finite square matrix A with entries in $\{0, 1\}$, the C^* -algebra $\mathcal{O}_{\mathfrak{L}^{\Lambda_A}}$ is canonically isomorphic to the Cuntz-Krieger algebra $\mathcal{O}_A$.

In this paper, we present a class of nonsofic subshifts $Z_N, N \in \mathbb{N}$ whose associated C^* -algebras $\mathcal{O}_{Z_N}$ are simple, purely infinite and not stably isomorphic to any Cuntz-Krieger algebra nor to Cuntz algebra. The subshifts $Z_N, N \in \mathbb{N}$ are coded systems whose languages are context free (cf. [2], [5], [13]). In studying a topological dynamical system, the topological entropy is very important quantity to measure complexity for the topological dynamical system. In [10], it was shown that the topological entropy for irreducible Markov shifts appear as the logarithm of the inverse temperature for admitting KMS-state for gauge action on the corresponding Cuntz-Krieger algebras. This result has been generalized to more general subshifts in [34] (cf. [22], [36]). Corresponding to these results, we will compute the topological entropy for the subshifts $Z_N, N \in \mathbb{N}$ and show that KMS-state for gauge action on the associated C^* -algebra $\mathcal{O}_{Z_N}$ exists if and only if the logarithm

of the inverse temperature is the topological entropy for the subshift Z_N , and the corresponding KMS-state is unique. We will prove

Theorem 1.1 (Theorem 3.7, Theorem 4.8 and Theorem 5.15).

- (i) For $N \in \mathbb{N}$, the C^* -algebra $\mathcal{O}_{Z_N}$ associated with the subshift Z_N is simple and purely infinite. It is the universal concrete C^* -algebra generated by two isometries T_1, T_2 and N partial isometries $S_i, i = 1, \dots, N$ satisfying the relations:

$$T_1 T_1^* + T_2 T_2^* + \sum_{j=1}^N S_j S_j^* = 1, \quad (1.4)$$

$$S_i^* S_i = 1 - \sum_{k=1}^{\infty} \sum_{m=0, k \neq 2m}^k \sum_{j=1}^N T_{2^m 1^{k-m}} S_j S_j^* T_{2^m 1^{k-m}}^*, \quad i = 1, \dots, N \quad (1.5)$$

where $T_{2^m 1^{k-m}}$ denotes $\underbrace{T_2 \cdots T_2}_{m \text{ times}} \underbrace{T_1 \cdots T_1}_{k-m \text{ times}}$. The infinite sum of the right hand side of the relation (1.5) is taken under strong operator topology on a Hilbert space.

- (ii) The K -groups for the C^* -algebra $\mathcal{O}_{Z_N}$ are as follows:

$$K_0(\mathcal{O}_{Z_N}) = \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z} \quad \text{and} \quad K_1(\mathcal{O}_{Z_N}) = 0.$$

The position of the unit $[1]$ in $K_0(\mathcal{O}_{Z_N}) = \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}$ is 0. Hence $\mathcal{O}_{Z_N}$ is not stably isomorphic to any Cuntz-Krieger algebra nor to the Cuntz algebra $\mathcal{O}_{\infty}$ for $N \geq 2$.

- (iii) There is a KMS-state for gauge action on $\mathcal{O}_{Z_N}$ if and only if the logarithm of the inverse temperature is $\log \beta_N$: the topological entropy of the subshift Z_N , where β_N is the unique solution β satisfying $\beta > N$ of the equation

$$\beta^6 - (N+3)\beta^5 + (3N+1)\beta^4 - 2(N-1)\beta^3 - (N+2)\beta^2 + N - 1 = 0. \quad (1.6)$$

The admitted KMS-state is unique.

The value β_N is increasing on N and satisfies $N < \beta_N < N+1$ such that

$$\lim_{N \rightarrow \infty} \frac{\beta_N}{N} = 1.$$

The class of the subshifts $Z_N, N \in \mathbb{N}$ is the first examples whose associated C^* -algebras are not stably isomorphic to any Cuntz-Krieger algebra nor to Cuntz algebra. For $N = 1$, the subshift Z_1 is nothing but the subshift Z named as the context free shift in [23, Example 1.2.9] and the associated C^* -algebra $\mathcal{O}_{Z_1}$ is the C^* -algebra $\mathcal{O}_Z$ studied in [26]. For other type of coded systems, see for example [21].

We will finally mention an application of our discussions to a classification problem in the theory of symbolic dynamical systems. By [30], [31], the K -groups $K_i(\mathcal{O}_{\Lambda})$ are invariant under not only topological conjugacy but also flow equivalence of subshifts Λ . Hence we know that the subshifts $Z_N, N \in \mathbb{N}$ are not flow equivalent to each other.

2. SUBSHIFTS AND THE C^* -ALGEBRAS ASSOCIATED WITH λ -GRAPH SYSTEMS

For a subshift Λ over alphabet Σ , we denote by $X_\Lambda \subset \Sigma^\mathbb{N}$ the set of all right-infinite sequences that appear in Λ . The topological dynamical system (X_Λ, σ) is called the one-sided subshift for Λ . A finite sequence $\mu = (\mu_1, \dots, \mu_k)$ of elements $\mu_j \in \Sigma$ is called a word. We denote by $|\mu|$ the length k of μ . A block $\mu = (\mu_1, \dots, \mu_k)$ is said to appear in $x = (x_i) \in \Sigma^\mathbb{Z}$ if $x_m = \mu_1, \dots, x_{m+k-1} = \mu_k$ for some $m \in \mathbb{Z}$. For a number $k \in \mathbb{N}$, let $B_k(\Lambda)$ be the set of all words of length k in $\Sigma^\mathbb{Z}$ appearing in some $x \in \Lambda$. Put $B_*(\Lambda) = \cup_{k=0}^\infty B_k(\Lambda)$ where $B_0(\Lambda)$ denotes the empty word $\emptyset$. Set

$$\Lambda_l(x) = \{\mu \in B_l(\Lambda) \mid \mu x \in X_\Lambda\} \quad \text{for } x \in X_\Lambda, \quad l \in \mathbb{Z}_+.$$

We define a nested sequence of equivalence relations in the space X_Λ as follows ([25], [27], [28]). Two points $x, y \in X_\Lambda$ are said to be l -past equivalent, written as $x \sim_l y$, if $\Lambda_l(x) = \Lambda_l(y)$. Denote by $\Omega_l = X_\Lambda / \sim_l$ the quotient space of X_Λ under $\sim_l$. For $x, y \in X_\Lambda$ and $\mu \in B_k(\Lambda)$, one sees that

- (i) if $x \sim_l y$, we have $x \sim_m y$ for $m < l$,
- (ii) if $x \sim_l y$ and $\mu x \in X_\Lambda$, we have $\mu y \in X_\Lambda$ and $\mu x \sim_{l-k} \mu y$ for $l > k$.

We have the following sequence of surjections in a natural way:

$$\Omega_0 \leftarrow \Omega_1 \leftarrow \Omega_2 \leftarrow \dots \leftarrow \Omega_l \leftarrow \Omega_{l+1} \leftarrow \dots,$$

where Ω_0 is a singleton. The subshift Λ is a sofic shift if and only if $\Omega_l = \Omega_{l+1}$ for some $l \in \mathbb{N}$. For a fixed $l \in \mathbb{Z}_+$, let $F_i^l, i = 1, 2, \dots, m(l)$ be the set of all l -past equivalence classes of X_Λ so that X_Λ is a disjoint union of $F_i^l, i = 1, 2, \dots, m(l)$. Then the canonical λ -graph system $\mathfrak{L}^\Lambda = (V^\Lambda, E^\Lambda, \lambda^\Lambda, \iota^\Lambda)$ for Λ is defined as follows ([28]). The vertex set V_l at level l consist of the sets $F_i^l, i = 1, \dots, m(l)$. We write an edge with label a from the vertex $F_i^l \in V_l$ to $F_j^{l+1} \in V_{l+1}$ if $ax \in F_j^{l+1}$ for some $x \in F_i^{l+1}$. We denote by $E_{l,l+1}$ the set of all edges from V_l to V_{l+1} . There exists a natural map $\iota_{l,l+1}^\Lambda$ from V_{l+1} to V_l by mapping F_j^{l+1} to F_i^l when F_i^l contains F_j^{l+1} . Set $V^\Lambda = \cup_{l=0}^\infty V_l$ and $E^\Lambda = \cup_{l=0}^\infty E_{l,l+1}$. The labeling of edges is denoted by $\lambda^\Lambda : E \rightarrow \Sigma$.

Let $\mathfrak{L} = (V, E, \lambda, \iota)$ be a λ -graph system over Σ . The C^* -algebra $\mathcal{O}_\mathfrak{L}$ associated with $\mathfrak{L}$ is the universal C^* -algebra generated by partial isometries $S_\alpha, \alpha \in \Sigma$ and projections $E_i^l, i = 1, 2, \dots, m(l), l \in \mathbb{Z}_+$ satisfying the following operator relations:

$$\sum_{\alpha \in \Sigma} S_\alpha S_\alpha^* = 1, \tag{2.1}$$

$$\sum_{i=1}^{m(l)} E_i^l = 1, \quad E_i^l = \sum_{j=1}^{m(l+1)} I_{l,l+1}(i, j) E_j^{l+1}, \tag{2.2}$$

$$S_\alpha S_\alpha^* E_i^l = E_i^l S_\alpha S_\alpha^*, \tag{2.3}$$

$$S_\alpha^* E_i^l S_\alpha = \sum_{j=1}^{m(l+1)} A_{l,l+1}(i, \alpha, j) E_j^{l+1}, \tag{2.4}$$

for $i = 1, 2, \dots, m(l), l \in \mathbb{Z}_+, \alpha \in \Sigma$, where $V_l = \{v_1^l, \dots, v_{m(l)}^l\}$,

$$A_{l,l+1}(i, \alpha, j) = \begin{cases} 1 & \text{if } s(e) = v_i^l, \lambda(e) = \alpha, t(e) = v_j^{l+1} \text{ for some } e \in E_{l,l+1}, \\ 0 & \text{otherwise,} \end{cases}$$

$$I_{l,l+1}(i, j) = \begin{cases} 1 & \text{if } \iota_{l,l+1}(v_j^{l+1}) = v_i^l, \\ 0 & \text{otherwise} \end{cases}$$

for $i = 1, 2, \dots, m(l), j = 1, 2, \dots, m(l+1), \alpha \in \Sigma$. If $\mathfrak{L}$ satisfies λ -condition (I) and is λ -irreducible, the C^* -algebra $\mathcal{O}_{\mathfrak{L}}$ is simple and purely infinite ([32], [33]). By the universality, the correspondence $\rho_z^{\mathfrak{L}} : z \in \mathbb{C}$ with $|z| = 1$ defined by $\rho_z^{\mathfrak{L}}(S_{\alpha}) = zS_{\alpha}, \rho_z^{\mathfrak{L}}(E_i^l) = E_i^l$ for $\alpha \in \Sigma, i = 1, \dots, m(l), l \in \mathbb{Z}_+$ yields an action $\rho^{\mathfrak{L}} : z \in \mathbb{T} \longrightarrow \rho_z^{\mathfrak{L}} \in \text{Aut}(\mathcal{O}_{\mathfrak{L}})$ called gauge action.

Let $\mathcal{A}_{\mathfrak{L},l}$ be the C^* -subalgebra of $\mathcal{O}_{\mathfrak{L}}$ generated by the projections $E_i^l, i = 1, \dots, m(l)$. We denote by $\mathcal{A}_{\mathfrak{L}}$ the C^* -subalgebra of $\mathcal{O}_{\mathfrak{L}}$ generated by the all projections $E_i^l, i = 1, \dots, m(l), l \in \mathbb{Z}_+$. Let $\lambda_{\mathfrak{L}}$ be the positive operator on the algebra $\mathcal{A}_{\mathfrak{L}}$ defined by

$$\lambda_{\mathfrak{L}}(X) = \sum_{\alpha \in \Sigma} S_{\alpha}^* X S_{\alpha} \quad \text{for } X \in \mathcal{A}_{\mathfrak{L}}.$$

Let $\lambda_l : \mathcal{A}_{\mathfrak{L},l} \longrightarrow \mathcal{A}_{\mathfrak{L},l+1}$ be the restriction of $\lambda_{\mathfrak{L}}$ to $\mathcal{A}_{\mathfrak{L},l}$ and $\iota_l : \mathcal{A}_{\mathfrak{L},l} \hookrightarrow \mathcal{A}_{\mathfrak{L},l+1}$ the natural inclusion.

We will in this paper study the C^* -algebras $\mathcal{O}_{\mathfrak{L}^{Z_N}}$ associated with the canonical λ -graph systems $\mathfrak{L}^{Z_N}$ for nonsofic subshifts $Z_N, N \in \mathbb{N}$ defined in the next section. The C^* -algebras $\mathcal{O}_{\mathfrak{L}^{Z_N}}$ will be denoted by $\mathcal{O}_{Z_N}$.

3. THE C^* -ALGEBRA $\mathcal{O}_{Z_N}$

For a finite set Σ , denote by Σ^* the set of all words of Σ . A (finite or infinite) collection $\mathcal{C}(\subset \Sigma^*)$ of words over Σ is said to be uniquely decipherable if whenever $\alpha_1 \alpha_2 \dots \alpha_n = \gamma_1 \gamma_2 \dots \gamma_m$ with $\alpha_i, \gamma_j \in \mathcal{C}$, then $n = m$ and $\alpha_i = \gamma_i$ for $i = 1, \dots, n$. A uniquely decipherable set $\mathcal{C}$ is called a code. Blanchard and Hansel [2] have introduced the notion of coded system. A subshift Λ is called a *coded system* if Λ is the closure of the set of biinfinite sequences obtained by freely concatenating the words in a code $\mathcal{C}$. It is denoted by $\Lambda_{\mathcal{C}}$. In this section, we will study certain coded systems written as $Z_N, N \in \mathbb{N}$. We fix a natural number $N \in \mathbb{N}$. Let Σ_N be a set $\{c, b, \alpha_1, \dots, \alpha_N\}$ of symbols. The subshift Z_N is defined to be the subshift over Σ_N whose forbidden words are

$$\mathcal{F}_N = \{\alpha_i b^m c^k \alpha_j \mid i, j = 1, \dots, N, m, k = 0, 1, \dots \text{ with } m \neq k\}$$

where the word $\alpha_i b^m c^k \alpha_j$ means $\alpha_i \underbrace{b \dots b}_{m \text{ times}} \underbrace{c \dots c}_{k \text{ times}} \alpha_j$. It is not a sofic subshift and hence not a Markov shift. Put

$$W_{b,c} = \{w \in \{b, c\}^* \mid w \neq b^m c^k \text{ for } m, k = 0, 1, \dots \text{ with } m \neq k\}$$

the set of all words of b, c that are not of the form $b^m c^k$ for $m, k = 0, 1, \dots$ with $m \neq k$.

Proposition 3.1. *The set $\mathcal{C}_N := \{\alpha_i w \mid i = 1, \dots, N, w \in W_{b,c}\}$ is a code such that $\Lambda_{\mathcal{C}_N} = Z_N$.*

Proof. It is clear that $\mathcal{C}_N$ is a code. The inclusion relation $\Lambda_{\mathcal{C}_N} \subset Z_N$ is obvious. Conversely, the forbidden words $\mathcal{F}_N$ are not admissible in $\Lambda_{\mathcal{C}_N}$. Hence $\Lambda_{\mathcal{C}_N} = Z_N$. $\square$

Define sequences of subsets of X_{Z_N} as in the following way.

$$P_0 = \{c^k b^\infty \mid k \geq 0\} \cup \{b^k c^m b y \in X_{Z_N} \mid k \geq 0, m \geq 1, y \in X_{Z_N}\}$$

and

$$E_n = \{c^n \alpha_j y \in X_{Z_N} \mid y \in X_{Z_N}, j = 1, \dots, N\},$$

$$Q_l = \bigcup_{n > l} E_n,$$

$$F_n = \{b^m c^{m+n} \alpha_j y \in X_{Z_N} \mid m \geq 1, y \in X_{Z_N}, j = 1, \dots, N\},$$

$$R_l = \{b^m c^k \alpha_j y \in X_{Z_N} \mid m \geq 1, k \geq 0, m+n \neq k \text{ for } n = 0, 1, \dots, l, j = 1, \dots, N\}$$

for $l, n = 0, 1, \dots$

Lemma 3.2. *For each $l \in \mathbb{N}$, the space X_{Z_N} is decomposed as a disjoint union:*

$$X_{Z_N} = P_0 \sqcup_{n=0}^{l-1} E_n \sqcup Q_{l-1} \sqcup_{n=0}^{l-1} F_n \sqcup R_{l-1}.$$

This decomposition of X_{Z_N} into $2l+3$ -components corresponds to the l -past equivalence classes of X_{Z_N} .

Irreducibility and condition (I) for square matrices with entries in $\{0, 1\}$ have been generalized to λ -graph systems as λ -irreducibility and λ -condition (I) respectively ([33]).

Lemma 3.3. *The canonical λ -graph system $\mathfrak{L}^{Z_N}$ satisfies λ -condition (I) and λ -irreducible.*

Proof. For any natural number l , set $L = l + 4$. For a word $\nu \in B_{l+2}(Z_N)$, put $\mu = \nu cb \in B_L(Z_N)$. Then we have for $x \in X_{Z_N}$

$$\begin{aligned} \mu x \in E_k & \quad \text{if} \quad \nu = c^k \alpha_j^{l-k+2} \text{ for some } j = 1, \dots, N \quad (k = 0, 1, \dots, l-1), \\ \mu x \in Q_{l-1} & \quad \text{if} \quad \nu = c^l \alpha_j^2 \text{ for some } j = 1, \dots, N, \\ \mu x \in F_k & \quad \text{if} \quad \nu = bc^{k+1} \alpha_j^{l-k} \text{ for some } j = 1, \dots, N \quad (k = 0, 1, \dots, l-1), \\ \mu x \in R_{l-1} & \quad \text{if} \quad \nu = b^2 c \alpha_j^{l-1} \text{ for some } j = 1, \dots, N, \\ \mu x \in P_0 & \quad \text{if} \quad \nu = cb^2. \end{aligned}$$

Since the family $E_0, F_0, E_1, F_1, \dots, E_{l-1}, F_{l-1}, Q_{l-1}, R_{l-1}, P_0$ represents the set of all l -past equivalence classes, the canonical λ -graph system $\mathfrak{L}^{Z_N}$ for the subshift Z_N is λ -irreducible. It is easy to see that $\mathfrak{L}^{Z_N}$ satisfies λ -condition (I). $\square$

Therefore we conclude by Lemma 3.3,

Corollary 3.4. *The C^* -algebra $\mathcal{O}_{Z_N}$ is simple and purely infinite.*

The C^* -algebra $\mathcal{O}_{Z_N}$ is generated by $N+2$ partial isometries $S_c, S_b, S_{\alpha_1}, \dots, S_{\alpha_N}$. We set $T_1 := S_c, T_2 := S_b, S_j := S_{\alpha_j}, j = 1, \dots, N$. Since for any $x \in X_{Z_N}$, both cx and bx are admissible and hence belong to X_{Z_N} , both the operators T_1 and T_2 are isometries. Since for any $x \in X_{Z_N}$, $\alpha_i x \in X_{Z_N}$ if and only if $\alpha_j x \in X_{Z_N}$ for $i, j = 1, \dots, N$, one has $S_i^* S_i = S_j^* S_j$ for $i, j = 1, \dots, N$. It has been proved in [29] that for a subshift Λ in general, the associated C^* -algebra $\mathcal{O}_\Lambda$ can be realized as a universal C^* -algebra as in the following way:

Lemma 3.5 ([29], cf. [4]). *For a subshift Λ over $\Sigma = \{1, 2, \dots, n\}$, the C^* -algebra $\mathcal{O}_\Lambda$ associated with Λ is the universal concrete C^* -algebra generated by n partial isometries $S_i, i = 1, 2, \dots, n$ subject to the following relations:*

- (i) $\sum_{j=1}^n S_j S_j^* = 1,$
- (ii) $S_i^* S_i = 1 - \sum_{k=1}^\infty \sum_{\nu \in L_i^k} S_\nu S_\nu^*, \quad i = 1, 2, \dots, n$

where $L_i^k = \{\nu_1 \cdots \nu_k \in B_k(\Lambda) \mid i\nu_1 \cdots \nu_{k-1} \in B_k(\Lambda), i\nu_1 \cdots \nu_{k-1}\nu_k \notin B_{k+1}(\Lambda)\}$. The infinite sum of the right hand side of the relation (ii) is taken under strong operator topology on a Hilbert space.

The above lemma means that there exists a representation of $\mathcal{O}_\Lambda$ in operators on a Hilbert space such that the canonical generators satisfy the relations both (i) and (ii). Conversely, if there exist n partial isometries on a Hilbert space satisfying the above relations, then there exists a canonical surjective homomorphism from $\mathcal{O}_\Lambda$ to the C^* -algebra generated by them.

We apply the preceding lemma to our C^* -algebra $\mathcal{O}_{Z_N}$. The following lemma is clear.

Lemma 3.6. $L_{\alpha_i}^k = \{b^m c^l \alpha_j \mid m + l = k - 1, m \neq l, \text{ and } j = 1, \dots, N\}$ for $i = 1, \dots, N$ and $k = 2, 3, \dots$.

Thus we obtain

Theorem 3.7. *For $N \in \mathbb{N}$, the C^* -algebra $\mathcal{O}_{Z_N}$ associated with the subshift Z_N is simple and purely infinite. It is the universal concrete C^* -algebra generated by two isometries T_1, T_2 and N partial isometries $S_j, j = 1, \dots, N$ satisfying the following relations:*

- (i) $T_1 T_1^* + T_2 T_2^* + \sum_{j=1}^N S_j S_j^* = 1,$
 - (ii) $S_i^* S_i = 1 - \sum_{k=1}^\infty \sum_{m=0, k \neq 2m}^k \sum_{j=1}^N T_{2^m 1^{k-m}} S_j S_j^* T_{2^m 1^{k-m}}^*, \quad i = 1, \dots, N$
- where $T_{2^m 1^{k-m}}$ denotes $\underbrace{T_2 \cdots T_2}_{m \text{ times}} \underbrace{T_1 \cdots T_1}_{k-m \text{ times}}$. The infinite sum of the right hand side of the relation (ii) is taken under strong operator topology on a Hilbert space.

4. THE K-THEORY FOR $\mathcal{O}_{Z_N}$

In this section, we compute the K-groups for $\mathcal{O}_{Z_N}$. Let $(M_{l,l+1}, I_{l,l+1})_{l \in \mathbb{Z}_+}$ be the nonnegative matrix system for the symbolic matrix system $(\mathcal{M}_{l,l+1}, I_{l,l+1})_{l \in \mathbb{Z}_+}$ of the canonical λ -graph system $\mathfrak{L}^{Z_N}$. Let $m(l)$ be the row size of the matrix $M_{l,l+1}$. The main tool is the following K-theory formulae proved in [25], [32]. The formulae hold for the C^* -algebras associated with λ -graph systems in general.

Lemma 4.1. ([25], [32])

- (i) $K_0(\mathcal{O}_{Z_N}) = \varinjlim \{\mathbb{Z}^{m(l+1)} / (M_{l,l+1}^t - I_{l,l+1}^t) \mathbb{Z}^{m(l)}, \overline{I}_{l,l+1}^t\}.$
 - (ii) $K_1(\mathcal{O}_{Z_N}) = \varinjlim \{\text{Ker}(M_{l,l+1}^t - I_{l,l+1}^t) \text{ in } \mathbb{Z}^{m(l)}, I_{l,l+1}^t\}.$
- where $M_{l,l+1}^t, I_{l,l+1}^t$ are the transposes of the matrices $M_{l,l+1}, I_{l,l+1}$ respectively, and $\overline{I}_{l,l+1}^t : \mathbb{Z}^{m(l)} / (M_{l-1,l}^t - I_{l-1,l}^t) \mathbb{Z}^{m(l-1)} \longrightarrow \mathbb{Z}^{m(l+1)} / (M_{l,l+1}^t - I_{l,l+1}^t) \mathbb{Z}^{m(l)}$ is the natural quotient map induced by $I_{l,l+1}^t$.

We will first find the symbolic matrix system $(\mathcal{M}_{l,l+1}, I_{l,l+1})_{l \in \mathbb{Z}_+}$ of the canonical λ -graph system $\mathfrak{L}^{Z_N}$. We have

- (1) $\alpha_i P_0 \subset P_0$ and $\alpha_i E_0, \alpha_i F_0 \subset E_0$ for $i = 1, \dots, N,$

- (2) $bP_0 \subset P_0$, $bE_0, bF_0, bQ_l, bR_l \subset R_{l-1}$ and $bE_n, bF_n \subset F_{n-1}$ for $n = 1, 2, \dots, l$,
(3) $cP_0, cF_n, cR_l \subset P_0$ for $n = 0, 1, \dots, l$, $cE_n \subset E_{n+1}$ for $n = 0, 1, \dots, l-2$ and $cE_{l-1}, cE_l, cQ_l \subset Q_{l-1}$,

and

$$Q_{l-1} = E_l \sqcup Q_l \quad \text{and} \quad R_{l-1} = F_l \sqcup R_l.$$

The we can represent the transposes $\mathcal{M}_{l,l+1}^t$ and $I_{l,l+1}^t$ of the matrices $\mathcal{M}_{l,l+1}$ and $I_{l,l+1}$ respectively as

$$\mathcal{M}_{l,l+1}^t = \begin{bmatrix} b+c & \alpha_1 + \dots + \alpha_N & 0 & \dots & \dots & \dots & \dots & \dots & 0 & 0 \\ 0 & \alpha_1 + \dots + \alpha_N & 0 & c & 0 & \dots & \dots & \dots & 0 & b \\ c & \alpha_1 + \dots + \alpha_N & 0 & \dots & \dots & \dots & \dots & \dots & 0 & b \\ 0 & 0 & b & 0 & 0 & c & 0 & \dots & 0 & 0 \\ c & 0 & b & 0 & \dots & \dots & \dots & \dots & 0 & 0 \\ 0 & 0 & 0 & 0 & b & 0 & 0 & c & 0 & 0 \\ c & 0 & 0 & 0 & b & 0 & \dots & \dots & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & b & 0 & 0 & c \\ c & 0 & 0 & 0 & 0 & 0 & b & 0 & \dots & 0 \\ \vdots & & & & \ddots & \ddots & \ddots & & \ddots & \vdots \\ \vdots & & & & & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & \dots & \dots & \dots & \dots & 0 & b & 0 & 0 & c & 0 \\ c & 0 & \dots & \dots & \dots & \dots & 0 & b & 0 & 0 & 0 & 0 \\ 0 & \dots & \dots & \dots & \dots & \dots & 0 & b & c & 0 & 0 & 0 \\ c & 0 & \dots & \dots & \dots & \dots & 0 & b & 0 & 0 & 0 & 0 \\ 0 & 0 & \dots & \dots & \dots & \dots & 0 & c & b & 0 & 0 & 0 \\ c & 0 & \dots & \dots & \dots & \dots & 0 & 0 & b & 0 & 0 & 0 \end{bmatrix},$$

and

$$I_{l,l+1}^t = \begin{bmatrix} 1 & 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ 0 & 1 & 0 & \dots & \dots & \dots & \dots & \dots & 0 \\ 0 & 0 & 1 & 0 & \dots & \dots & \dots & \dots & 0 \\ 0 & 0 & 0 & 1 & 0 & \dots & \dots & \dots & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & \dots & \dots & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ \vdots & & & & \ddots & \ddots & \ddots & & \ddots & \vdots & \vdots \\ \vdots & & & & & \ddots & \ddots & \ddots & \ddots & \vdots & \vdots \\ \vdots & & & & & & \ddots & \ddots & \ddots & \vdots & \vdots \\ 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & 0 & 0 \\ 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & 0 & 0 \\ 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & 0 & 0 \\ 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

along the ordered basis

$$\begin{aligned} & (P_0, E_0, F_0, E_1, F_1, \dots, E_l, F_l, Q_l, R_l) - \text{row} \\ & (P_0, E_0, F_0, E_1, F_1, \dots, E_{l-1}, F_{l-1}, Q_{l-1}, R_{l-1}) - \text{column.} \end{aligned}$$

The commutative C^* -algebras $\mathcal{A}_{\mathcal{L}^{Z_N}, l}$ and $\mathcal{A}_{\mathcal{L}^{Z_N}}$ are denoted by $\mathcal{A}_{Z_N, l}$ and $\mathcal{A}_{Z_N}$ respectively. The set of the minimal projections $\mathcal{A}_{Z_N, l}$ correspond to the set of all l -past equivalence classes $P_0, E_0, F_0, E_1, F_1, \dots, E_{l-1}, F_{l-1}, Q_{l-1}, R_{l-1}$ which are denoted by $p_0, e_0, f_0, e_1, f_1, \dots, e_{l-1}, f_{l-1}, q_{l-1}, r_{l-1}$ respectively. Hence $\dim(\mathcal{A}_l) = m(l) = 2l+3$ for $l \geq 2$. We can then represent the induced matrix on the K_0 -groups

$$M_{l, l+1}^t (= \lambda_{l*}) : K_0(\mathcal{A}_{Z_N, l}) = \mathbb{Z}^{2l+3} \rightarrow K_0(\mathcal{A}_{Z_N, l+1}) = \mathbb{Z}^{2l+5}$$

of the operator λ_l to be the matrix $M_{l, l+1}^t$ as follows:

$$M_{l, l+1}^t = \begin{bmatrix} 2 & N & 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 & 0 \\ 0 & N & 0 & 1 & 0 & \dots & \dots & \dots & \dots & 0 & 1 \\ 1 & N & 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & \dots & \dots & 0 & \\ 1 & 0 & 1 & 0 & \dots & \dots & \dots & \dots & \dots & 0 & \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & \dots & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & \dots & \dots & \dots & 0 & \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & \dots & \dots & 0 \\ \vdots & & & & & \ddots & \ddots & \ddots & & \ddots & \vdots \\ \vdots & & & & & & \ddots & \ddots & \ddots & & \vdots \\ \vdots & & & & & & & \ddots & \ddots & & \vdots \\ 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & 0 \\ 1 & 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & 0 \\ 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 1 & 0 \\ 1 & 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & 0 \\ 0 & 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 1 & \\ 1 & 0 & \dots & \dots & \dots & \dots & \dots & 0 & 0 & 1 & \end{bmatrix}.$$

The natural inclusion ι_l from $\mathcal{A}_{Z_N, l}$ to $\mathcal{A}_{Z_N, l+1}$ induces the matrix

$$I_{l, l+1}^t (= \iota_{l*}) : K_0(\mathcal{A}_{Z_N, l}) = \mathbb{Z}^{2l+3} \rightarrow K_0(\mathcal{A}_{Z_N, l+1}) = \mathbb{Z}^{2l+5}.$$

As

$$M_{l,l+1}^t - I_{l,l+1}^t = \begin{bmatrix} 1 & N & 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 & 0 \\ 0 & N-1 & 0 & 1 & 0 & \dots & \dots & \dots & \dots & 0 & 1 \\ 1 & N & -1 & 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 \\ 0 & 0 & 1 & -1 & 0 & 1 & 0 & \dots & \dots & 0 & 0 \\ 1 & 0 & 1 & 0 & -1 & 0 & \dots & \dots & \dots & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 & 1 & 0 & \dots & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & -1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & -1 & 0 & \dots \\ \vdots & & & & & \ddots & \ddots & \ddots & & \ddots & \vdots \\ \vdots & & & & & & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 & 1 & -1 & 0 & 1 & 0 \\ 1 & 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & \dots & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & \dots & \dots & \dots & \dots & \dots & 0 & 1 & 0 & -1 \\ 0 & 0 & \dots & \dots & \dots & \dots & \dots & 0 & 0 & 1 \\ 1 & 0 & \dots & \dots & \dots & \dots & \dots & \dots & \dots & 0 \end{bmatrix},$$

we easily see that

Lemma 4.2. $\text{Ker}(M_{l,l+1}^t - I_{l,l+1}^t) = 0$ for $2 \leq l \in \mathbb{N}$.

Thus we have by Lemma 4.1,

Proposition 4.3. $K_1(\mathcal{O}_{Z_N}) \cong 0$.

We will next compute $K_0(\mathcal{O}_{Z_N})$. For an integer n , we denote by $q(n) \in \mathbb{Z}$ the quotient of n by N and by $r(n) \in \{0, 1, \dots, N-1\}$ its residue such as $n = q(n)N + r(n)$.

Lemma 4.4. Fix $l = 2, 3, \dots$. For $z = \begin{bmatrix} z_1 \\ \vdots \\ z_{2l+5} \end{bmatrix} \in \mathbb{Z}^{2l+5}$, put inductively

$$\begin{aligned} x_{2l+3} &= z_{2l+4}, \\ x_{2l+1} &= z_{2l+2}, \\ x_{2l-1} &= z_{2l+1} - z_{2l+5} + z_{2l+2}, \\ x_{2l-3} &= z_{2l-1} - z_{2l+5} + x_{2l-1}, \\ x_{2l-5} &= z_{2l-3} - z_{2l+5} + x_{2l-3}, \\ x_{2l-7} &= z_{2l-5} - z_{2l+5} + x_{2l-5}, \\ &\vdots \\ x_3 &= z_5 - z_{2l+5} + x_5, \\ x_1 &= z_{2l+5}, \end{aligned}$$

and

$$\begin{aligned}
x_2 &= q(z_1 - z_{2l+5}), \\
x_4 &= z_2 - z_{2l+4} - (N-1)x_2, \\
x_6 &= z_4 - x_3 + x_4, \\
x_8 &= z_6 - x_5 + x_6, \\
&\vdots \\
x_{2l} &= z_{2l-2} - x_{2l-3} + x_{2l-2}, \\
x_{2l+2} &= z_{2l} - x_{2l-1} + x_{2l}.
\end{aligned}$$

Set

$$\begin{aligned}
r_{l+1}(z) &= r(z_1 - z_{2l+5}) \in \{0, 1, \dots, N-1\}, \\
\eta_{l+1}(z) &= z_3 - (x_1 + Nx_2) + x_3 - x_{2l+3} \\
&= -z_1 + (z_3 + z_5 + \dots + z_{2l+1}) + z_{2l+2} - z_{2l+4} - (l-1)z_{2l+5} + r_{l+1}(z), \\
\psi_{l+1}(z) &= z_{2l+3} - x_1 - x_{2l+1} + x_{2l+3} \\
&= -z_{2l+2} + z_{2l+3} + z_{2l+4} - z_{2l+5}.
\end{aligned}$$

Then we have

$$\begin{bmatrix} z_1 \\ \vdots \\ z_{2l+5} \end{bmatrix} = (M_{l,l+1}^t - I_{l,l+1}^t) \begin{bmatrix} x_1 \\ \vdots \\ x_{2l+3} \end{bmatrix} + \begin{bmatrix} r_{l+1}(z) \\ 0 \\ \eta_{l+1}(z) \\ 0 \\ \vdots \\ 0 \\ \psi_{l+1}(z) \\ 0 \\ 0 \end{bmatrix}.$$

Proof. For $[y_i]_{i=1}^{2l+3} \in \mathbb{Z}^{2l+3}$, one sees

$$(M_{l,l+1}^t - I_{l,l+1}^t) \begin{bmatrix} y_1 \\ \vdots \\ y_{2l+3} \end{bmatrix} = \begin{bmatrix} y_1 + Ny_2 \\ (N-1)y_2 + y_4 + y_{2l+3} \\ y_1 + Ny_2 - y_3 + y_{2l+3} \\ y_3 - y_4 + y_6 \\ y_1 + y_3 - y_5 \\ \vdots \\ y_{2k-1} - y_{2k} + y_{2k+2} \\ y_1 + y_{2k-1} - y_{2k+1} \\ \vdots \\ y_{2l-1} - y_{2l} + y_{2l+2} \\ y_1 + y_{2l-1} - y_{2l+1} \\ y_{2l+1} \\ y_1 + y_{2l+1} - y_{2l+3} \\ y_{2l+3} \\ y_1 \end{bmatrix}.$$

Hence the assertion is straightforward. $\square$

Lemma 4.5. For $z = [z_i]_{i=1}^{2l+5} \in \mathbb{Z}^{2l+5}$, one has

$$r_{l+1}(z) = 0 \text{ in } \{0, 1, \dots, N-1\} \quad \text{and} \quad \eta_{l+1}(z) = \psi_{l+1}(z) = 0 \text{ in } \mathbb{Z}$$

if and only if there exists $y = [y_i]_{i=1}^{2l+3} \in \mathbb{Z}^{2l+3}$ such that $z = (M_{l,l+1}^t - I_{l,l+1}^t)(y)$.

Proof. The only if part follows from the preceding lemma. We will show the if part. Suppose that there exists $y = [y_i]_{i=1}^{2l+3} \in \mathbb{Z}^{2l+3}$ such that $z = (M_{l,l+1}^t - I_{l,l+1}^t)(y)$. Define $[x_i]_{i=1}^{2l+3} \in \mathbb{Z}^{2l+3}$ for z as in the preceding lemma and put

$$\rho_{l+1}(z) = [r_{l+1}(z), 0, \eta_{l+1}(z), 0, \dots, 0, \psi_{l+1}(z), 0, 0]^t \in \mathbb{Z}^{2l+5}.$$

Put $w_i = y_i - x_i$, $1 \leq i \leq 2l+3$ so that the equality

$$(M_{l,l+1}^t - I_{l,l+1}^t)([w_i]_{i=1}^{2l+3}) = \rho_{l+1}(z) \quad (4.1)$$

holds. From the $2l+5$ -th row in the equality (4.1), one sees that $w_1 = 0$. From the $2l+4$ -th row and the $2l+2$ -th row, one sees that $w_{2l+3} = w_{2l+1} = 0$ so that $\psi_{l+1}(z) = 0$. Inductively, from the $2k+1$ -row, one sees that $w_{2k-1} = w_{2k+1} - w_1 = 0$ for $k = l, l-1, \dots, 2$. From the first row with $w_1 = 0$, one has $Nw_2 = r_{l+1}(z) \in \{0, 1, \dots, N-1\}$ so that $w_2 = 0$ and $r_{l+1}(z) = 0$. Hence one has from the third row $\eta_{l+1}(z) = w_1 + Nw_2 - w_3 + w_{2l+3} = 0$. $\square$

Lemma 4.6. The map $\xi_{l+1} : [z_i]_{i=1}^{2l+5} \in \mathbb{Z}^{2l+5} \longrightarrow (r_{l+1}(z), \eta_{l+1}(z), \psi_{l+1}(z)) \in \{0, 1, \dots, N-1\} \oplus \mathbb{Z} \oplus \mathbb{Z}$ induces an isomorphism from $\mathbb{Z}^{2l+5}/(M_{l,l+1}^t - I_{l,l+1}^t)\mathbb{Z}^{2l+3}$ onto $\mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$.

Proof. It suffices to show the surjectivity of ξ_{l+1} . For $(g, m, k) \in \{0, 1, \dots, N-1\} \oplus \mathbb{Z} \oplus \mathbb{Z}$, put $z = [g, 0, m, 0, \dots, 0, k, 0, 0]^t \in \mathbb{Z}^{2l+5}$. One then sees that

$$r_{l+1}(z) = g, \quad \eta_{l+1}(z) = m, \quad \psi_{l+1}(z) = k.$$

$\square$

We denote by $\bar{\xi}_{l+1}$ the above isomorphism from $\mathbb{Z}^{2l+5}/(M_{l,l+1}^t - I_{l,l+1}^t)\mathbb{Z}^{2l+3}$ onto $\mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}$ induced by ξ_{l+1} .

Lemma 4.7. The diagram

$$\begin{array}{ccc} \mathbb{Z}^{2l+3}/(M_{l-1,l}^t - I_{l-1,l}^t)\mathbb{Z}^{2l+1} & \xrightarrow{\overline{I}^{l,l+1}} & \mathbb{Z}^{2l+5}/(M_{l,l+1}^t - I_{l,l+1}^t)\mathbb{Z}^{2l+3} \\ \xi_l \downarrow & & \bar{\xi}_{l+1} \downarrow \\ \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} & \xrightarrow{L} & \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} \end{array}$$

is commutative, where $L = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$.

Proof. For $(g, m, k) \in \{0, 1, \dots, N-1\} \oplus \mathbb{Z} \oplus \mathbb{Z}$, put $z = [g, 0, m, 0, \dots, 0, k, 0, 0]^t \in \mathbb{Z}^{2l+5}$. Since $I_{l,l+1}^t(z) = [g, 0, m, 0, \dots, 0, k, 0, 0, 0]^t \in \mathbb{Z}^{2l+5}$, one sees that

$$r_{l+1}(I_{l,l+1}^t(z)) = g, \quad \eta_{l+1}(I_{l,l+1}^t(z)) = m + k, \quad \psi_{l+1}(I_{l,l+1}^t(z)) = 0.$$

Hence the above diagram is commutative. $\square$

Therefore we conclude

Theorem 4.8. $K_0(\mathcal{O}_{Z_N}) \cong \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}$, and the class $[1]$ of the unit 1 of the algebra $\mathcal{O}_{Z_N}$ in $K_0(\mathcal{O}_{Z_N})$ is $0 \oplus 0$ in $\mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}$.

Proof. By Lemma 4.1, it follows that

$$\begin{aligned} K_0(\mathcal{O}_{Z_N}) &= \varinjlim \{ \mathbb{Z}^{2l+5} / (M_{l,l+1}^t - I_{l,l+1}^t) \mathbb{Z}^{2l+3}, \overline{I}_{l,l+1}^t \} \\ &= \varinjlim \{ \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}, L \} \\ &\cong \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}. \end{aligned}$$

The class of the unit $[1]$ in $K_0(\mathcal{O}_{Z_N})$ corresponds to $[1]$ in $K_0(\mathcal{A}_{\mathfrak{L},l})$ that is represented to be the vector $[1, \dots, 1] \in \mathbb{Z}^{2l+3}$. Since $r_{l+1}([1]) = \eta_{l+1}([1]) = \psi_{l+1}([1]) = 0$, we see $\xi_{l+1}([1]) = (0, 0, 0)$. Thus the projection $[1]$ in $K_0(\mathcal{O}_{Z_N})$ represents $0 \oplus 0$ in $\mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}$. $\square$

By [8], we know $K_0(\mathcal{O}_\infty) = \mathbb{Z}$, $K_1(\mathcal{O}_\infty) = 0$ and the position of $[1]$ in $K_0(\mathcal{O}_\infty)$ is 1 in $\mathbb{Z}$. The classification theorem of purely infinite simple C^* -algebras proved by Kirchberg [15] and Philips [37], we have

Corollary 4.9. (i) *The C^* -algebra $\mathcal{O}_{Z_N}$ for $N \geq 2$ is not stably isomorphic to any Cuntz-Krieger algebra nor to the Cuntz algebra $\mathcal{O}_\infty$ of order infinity.*
(ii) *The C^* -algebra $\mathcal{O}_{Z_N}$ for $N = 1$ is not isomorphic to any Cuntz-Krieger algebra nor to the Cuntz algebra $\mathcal{O}_\infty$ of order infinity but stably isomorphic to $\mathcal{O}_\infty$. In fact, $\mathcal{O}_{Z_1}$ is isomorphic to $(\mathcal{O}_\infty)_{1-s_1s_1^*}$ where $\mathcal{O}_\infty$ is generated by a sequence $s_i, i = 1, 2, \dots$ of isometries with mutually orthogonal ranges.*

5. KMS-STATES FOR GAUGE ACTION AND TOPOLOGICAL ENTROPY

In this section, we study KMS-state for gauge action on $\mathcal{O}_{Z_N}$ and compute topological entropy for the subshift Z_N .

In studying KMS-state for gauge action on the C^* -algebras associated with subshifts, the lemma below proved in [34] plays a crucial rôle (cf. [22], [36]).

Lemma 5.1 ([34]). *For a λ -graph system $\mathfrak{L}$, if a state φ on $\mathcal{O}_{\mathfrak{L}}$ is a KMS-state for gauge action on $\mathcal{O}_{\mathfrak{L}}$ at the inverse temperature $\log \beta$ for some $1 < \beta \in \mathbb{R}$, it satisfies the condition $\varphi \circ \lambda_{\mathfrak{L}} = \beta \varphi$ on $\mathcal{A}_{\mathfrak{L}}$. Conversely, a state φ on $\mathcal{A}_{\mathfrak{L}}$ satisfying the condition $\varphi \circ \lambda_{\mathfrak{L}} = \beta \varphi$ can be uniquely extended to a KMS-state for gauge action on $\mathcal{O}_{\mathfrak{L}}$.*

We will apply this lemma to our C^* -algebra $\mathcal{O}_{Z_N}$ for the subshift Z_N . Let us find a state on $\mathcal{A}_{Z_N}$ satisfying the condition $\varphi \circ \lambda_{Z_N} = \beta \varphi$ for some real number β with $1 < \beta \in \mathbb{R}$, where λ_{Z_N} denotes $\lambda_{\mathfrak{L}^{Z_N}}$.

Let $\hat{p}_0, \hat{e}_0, \hat{f}_0, \hat{e}_1, \hat{f}_1, \dots, \hat{e}_l, \hat{f}_l, \hat{q}_l, \hat{r}_l$ be real numbers for $l \in \mathbb{N}$ satisfying the condition

$$\hat{p}_0 + \sum_{j=0}^l (\hat{e}_j + \hat{f}_j) + \hat{q}_l + \hat{r}_l = 1. \quad (5.1)$$

We consider the following equations :

$$M_{l,l+1}^t \begin{bmatrix} \hat{p}_0 \\ \hat{e}_0 \\ \hat{f}_0 \\ \hat{e}_1 \\ \hat{f}_1 \\ \hat{e}_2 \\ \hat{f}_2 \\ \vdots \\ \hat{e}_{l-1} \\ \hat{f}_{l-1} \\ \hat{e}_l \\ \hat{f}_l \\ \hat{q}_l \\ \hat{r}_l \end{bmatrix} = \beta \begin{bmatrix} \hat{p}_0 \\ \hat{e}_0 \\ \hat{f}_0 \\ \hat{e}_1 \\ \hat{f}_1 \\ \hat{e}_2 \\ \hat{f}_2 \\ \vdots \\ \hat{e}_{l-1} \\ \hat{f}_{l-1} \\ \hat{q}_{l-1} \\ \hat{r}_{l-1} \end{bmatrix} \quad \text{and} \quad I_{l,l+1}^t \begin{bmatrix} \hat{p}_0 \\ \hat{e}_0 \\ \hat{f}_0 \\ \hat{e}_1 \\ \hat{f}_1 \\ \hat{e}_2 \\ \hat{f}_2 \\ \vdots \\ \hat{e}_{l-1} \\ \hat{f}_{l-1} \\ \hat{e}_l \\ \hat{f}_l \\ \hat{q}_l \\ \hat{r}_l \end{bmatrix} = \begin{bmatrix} \hat{p}_0 \\ \hat{e}_0 \\ \hat{f}_0 \\ \hat{e}_1 \\ \hat{f}_1 \\ \hat{e}_2 \\ \hat{f}_2 \\ \vdots \\ \hat{e}_{l-1} \\ \hat{f}_{l-1} \\ \hat{q}_{l-1} \\ \hat{r}_{l-1} \end{bmatrix}$$

for a real number $\beta > 1$. This means that

$$2\hat{p}_0 + \sum_{i=0}^l \hat{f}_i + \hat{r}_l = \beta\hat{p}_0, \quad (5.2)$$

$$N(\hat{p}_0 + \hat{e}_0 + \hat{f}_0) = \beta\hat{e}_0, \quad (5.3)$$

$$\hat{e}_n + \hat{f}_n = \beta\hat{f}_{n-1}, \quad n = 1, 2, \dots, l, \quad (5.4)$$

$$\hat{e}_{n-1} = \beta\hat{e}_n, \quad n = 1, 2, \dots, l-1, \quad (5.5)$$

$$\hat{e}_{l-1} + \hat{e}_l + \hat{q}_l = \beta\hat{q}_{l-1}, \quad (5.6)$$

$$\hat{e}_0 + \hat{f}_0 + \hat{q}_l + \hat{r}_l = \beta\hat{r}_{l-1} \quad (5.7)$$

and the equations

$$\hat{e}_l + \hat{q}_l = \hat{q}_{l-1}, \quad (5.8)$$

$$\hat{f}_l + \hat{r}_l = \hat{r}_{l-1}. \quad (5.9)$$

These equations can be solved as in the following way by straightforward calculation. Put

$$g_N(\beta) = (\beta - 1)\beta\{(2N - 1)\beta - (N + 1)\},$$

$$h_N(\beta) = (\beta + 1)\{(\beta - N)(\beta - 2)(\beta - 1)^2\beta + (N - 2)(\beta - 1)\beta - N(\beta - 1) - 1\}$$

and

$$F_N(\beta) = h_N(\beta) - g_N(\beta).$$

Lemma 5.2.

$$\hat{p}_0 = \frac{1}{N} \{(\beta - N)\hat{e}_0 - N\hat{f}_0\}, \quad (5.10)$$

$$(\beta - 1)(\beta + 1)g_N(\beta)\hat{f}_0 = h_N(\beta)\hat{e}_0, \quad (5.11)$$

$$\hat{e}_n = \frac{\beta - 2}{\beta^{n+1}}, \quad (5.12)$$

$$\hat{f}_n = \beta^n \hat{f}_0 - (\beta^{n-1}\hat{e}_1 + \beta^{n-2}\hat{e}_2 + \cdots + \beta\hat{e}_{n-1} + \hat{e}_n), \quad (5.13)$$

$$\hat{q}_n = \frac{\beta - 2}{\beta^{n+1}(\beta - 1)}, \quad (5.14)$$

$$\hat{r}_n = (\beta - 2)\hat{p}_0 - (\hat{f}_0 + \hat{f}_1 + \cdots + \hat{f}_n), \quad n = 0, 1, 2, \dots, l. \quad (5.15)$$

We henceforth assume that the real numbers $\hat{p}_0, \hat{e}_0, \hat{f}_0, \hat{e}_1, \hat{f}_1, \dots, \hat{e}_l, \hat{f}_l, \hat{q}_l, \hat{r}_l$ are all nonnegative for all $l \in \mathbb{N}$.

Lemma 5.3. (i) $\beta > 2$ and $\hat{e}_n, \hat{q}_n > 0$ for $n = 0, 1, \dots, l$.

(ii) $\beta > N$.

(iii) $g_N(\beta) > 0$.

Proof. (i) Since $\hat{e}_1 \geq 0$, one sees that $\beta \geq 2$ by (5.12). Suppose that $\beta = 2$. One has $\hat{e}_n = \hat{q}_n = 0$ for $n = 0, 1, \dots, l$. (5.15) implies that $\hat{r}_n = \hat{f}_n = 0$ for $n = 0, 1, \dots, l$. It follows that $\hat{p}_0 = 0$ by (5.10), a contradiction to (5.1). Hence $\beta > 2$ and $\hat{e}_n, \hat{q}_n > 0$ for $n = 0, 1, \dots, l$.

(ii) Suppose next that $\beta \leq N$. (5.10) implies that $\hat{p}_0 = 0, \hat{f}_0 = 0$. Then (5.13) implies that $\hat{f}_n < 0$ a contradiction.

(iii) As

$$\frac{g_N(\beta)}{(\beta - 1)\beta} = (2N - 1)\beta - (N + 1) > (2N - 1)N - (N + 1) = (N - 1)^2 + N^2 - 2. \quad (5.16)$$

one sees that $g_N(\beta) > 0$ for $N \geq 2$ and $g_1(\beta) = (\beta - 2)(\beta - 1)\beta > 0$ $\square$

Lemma 5.4. The conditions $\hat{f}_n \geq 0$ for all $n = 0, 1, \dots$ imply $F_N(\beta) \geq 0$.

Proof. (5.12) and (5.13) imply

$$\hat{f}_n = \beta^n \left\{ \hat{f}_0 - \frac{1}{\beta^2 - 1} \left(1 - \frac{1}{\beta^{2n}} \right) \hat{e}_0 \right\} \quad (5.17)$$

so that the conditions $\hat{f}_n \geq 0$ for all $n = 0, 1, \dots$ imply

$$(\beta^2 - 1)\hat{f}_0 - \hat{e}_0 \geq -\frac{1}{\beta^{2n}}\hat{e}_0 \quad \text{for all } n = 0, 1, \dots \quad (5.18)$$

As $g_N(\beta) > 0$, by (5.11), the above equalities imply the inequality $h_N(\beta) - g_N(\beta) \geq 0$. $\square$

Lemma 5.5. The conditions $\hat{r}_n \geq 0$ for all $n = 0, 1, \dots$ imply $F_N(\beta) \leq 0$.

Proof. By (5.12) and (5.13), one sees that

$$\sum_{k=0}^n \hat{f}_k = \frac{\beta^{n+1} - 1}{\beta - 1} \hat{f}_0 + \left\{ \frac{1}{(\beta - 1)^2} - \frac{1}{(\beta - 1)^2(\beta + 1)} \right\} (\beta^{n+1} + \frac{1}{\beta^n}) \hat{e}_0.$$

(5.15) implies that

$$\begin{aligned}
& (\beta - 1)^2(\beta + 1)g_N(\beta)\hat{r}_n \\
&= [\frac{\beta - N}{N}(\beta - 2)(\beta - 1)^2(\beta + 1) - \{(\beta + 1) - (\beta^{n+1} + \frac{1}{\beta^n})\}]g_N(\beta)\hat{e}_0 \\
&\quad - [(\beta - 2)(\beta - 1)^2(\beta + 1) + (\beta - 1)(\beta + 1)(\beta^{n+1} - 1)]g_N(\beta)\hat{f}_0 \\
&= [\frac{\beta - N}{N}(\beta - 2)(\beta - 1)^2(\beta + 1) - \{(\beta + 1) - (\beta^{n+1} + \frac{1}{\beta^n})\}]g_N(\beta)\hat{e}_0 \\
&\quad - [(\beta - 2)(\beta - 1) + (\beta^{n+1} - 1)]h_N(\beta)\hat{e}_0 \\
&= -F_N(\beta)\beta^{n+1}\hat{e}_0 \\
&\quad + [\frac{\beta - N}{N}(\beta - 2)(\beta - 1)^2(\beta + 1) - \{(\beta + 1) - \frac{1}{\beta^n}\}]g_N(\beta)\hat{e}_0 \\
&\quad - [(\beta - 2)(\beta - 1) - 1]h_N(\beta)\hat{e}_0
\end{aligned}$$

Since

$$\liminf_{n \rightarrow \infty} \frac{(\beta - 1)^2(\beta + 1)g_N(\beta)\hat{r}_n}{\beta^{n+1}} = \liminf_{n \rightarrow \infty} \frac{(\beta - 1)^2(\beta + 1)g_N(\beta)\hat{r}_n}{\beta^{n+1}} = -F_N(\beta)\hat{e}_0 \geq 0$$

and $\hat{e}_0 > 0$, we see $F_N(\beta) \leq 0$. $\square$

Therefore we have

Corollary 5.6. *Suppose that $\hat{p}_0, \hat{e}_n, \hat{f}_n, \hat{q}_n, \hat{r}_n$ are all nonnegative for all $n = 0, 1, \dots$. Then $\beta > N$ and $F_N(\beta) = 0$.*

Conversely, for $N \geq 2$, we have

Lemma 5.7. *For $\beta > N$, $F_N(\beta) = 0$ implies $\hat{f}_n, \hat{r}_n > 0$ for all $n = 0, 1, \dots$.*

Proof. Assume that $F_N(\beta) = 0$. As $\beta > N$, (5.16) implies $g_N(\beta) > 0$. Since $h_N(\beta) = g_N(\beta)$, the inequality (5.18) hold, which means $\hat{f}_n > 0$ for all $n = 0, 1, \dots$ by (5.17).

We will next show that $\hat{r}_n > 0$ for all $n = 0, 1, \dots$. By the equality $g_N(\beta) = h_N(\beta)$, one has

$$\begin{aligned}
& (\beta - 1)^2(\beta + 1)g_N(\beta)\hat{r}_n \\
&= [\frac{\beta - N}{N}(\beta - 2)(\beta - 1)^2(\beta + 1) - \{(\beta + 1) - \frac{1}{\beta^n}\}]g_N(\beta)\hat{e}_0 \\
&\quad - [(\beta - 2)(\beta - 1) - 1]g_N(\beta)\hat{e}_0.
\end{aligned}$$

It follows that

$$\begin{aligned}
& [\frac{\beta - N}{N}(\beta - 2)(\beta - 1)^2(\beta + 1) - (\beta + 1)] - [(\beta - 2)(\beta - 1) - 1] \\
&= \frac{1}{N\beta}(\beta^3 - N + 1) > 0
\end{aligned}$$

so that $(\beta - 1)^2(\beta + 1)g_N(\beta)\hat{r}_n$ is positive. As $g_N(\beta) > 0$, one sees $\hat{r}_n > 0$ for all $n = 0, 1, \dots$. $\square$

Therefore we have

Corollary 5.8. *For $\beta > N$, $F_N(\beta) = 0$ if and only if $\hat{p}_0, \hat{e}_n, \hat{f}_n, \hat{q}_n, \hat{r}_n$ are all nonnegative for all $n = 0, 1, \dots$. In this case, they are all positive.*

Proof. For $N = 1$, the statement has been shown in [26]. $\square$

We note that the identity

$$F_N(\beta) = \beta^6 - (N+3)\beta^5 + (3N+1)\beta^4 - 2(N-1)\beta^3 - (N+2)\beta^2 + N - 1$$

holds. For $N = 1$, the above identity goes to

$$F_1(\beta) = \beta^6 - 4\beta^5 + 4\beta^4 - 3\beta^2.$$

The unique positive solution of $F_1(\beta) = 0$ is $\beta = 1 + \sqrt{1 + \sqrt{3}} = 2.652891 \dots$ ([26]).

We will next study positive solutions of the equations

$$F_N(\beta) = 0 \quad \text{for} \quad N \geq 2.$$

Lemma 5.9.

- (i) $F_2(3) < 0 < F_2(4)$.
- (ii) $F_N(N) < 0 < F_N(N+1)$ for $N \geq 3$.

Proof. By the identities

$$F_N(N) = -(N^2 + 1)(N^2 - N + 1),$$

$$F_N(N+1) = \{N(N-2)(N+1) - 1\}(N+1)^2 + N - 1$$

one sees that (iii) holds. $\square$

The following lemma is straightforward.

Lemma 5.10.

- (i) For $N = 2$, $F'_2(\beta) > 0$ for all $\beta \geq 3$. Hence $F_2(\beta) = 0$ has a unique solution in $[3, \infty)$.
- (ii) For $N = 3$, $F'_3(3) < 0$ and $F''_3(\beta) > 0$ for all $\beta \geq 3$. Hence $F_3(\beta) = 0$ has a unique solution in $[3, \infty)$.
- (iii) For $N \geq 4$, $F'_N(\beta) > 0$ for all $\beta \geq N$. Hence $F_N(\beta) = 0$ has a unique solution in $[N, \infty)$.

Therefore we have

Proposition 5.11. For $N \geq 1$, the equation $F_N(\beta) = 0$ has a unique solution in $[N, \infty)$. It satisfies $N < \beta < N+1$ for $N \geq 3$.

Lemma 5.12. For the unique $\beta > N$ satisfying $F_N(\beta) = 0$, one has

$$\begin{aligned} \hat{p}_0 &= \frac{(\beta-2)(\beta^2 - N\beta - 1)}{N(\beta^2 - 1)}, \\ \hat{e}_n &= \frac{\beta-2}{\beta^{n+1}}, \\ \hat{f}_n &= \frac{\beta-2}{(\beta^2-1)\beta^{n+1}}, \\ \hat{q}_n &= \frac{\beta-2}{\beta^{n+1}(\beta-1)}, \\ \hat{r}_n &= \frac{\beta-2}{N(\beta-1)^2(\beta+1)} \{(\beta-2)(\beta-1)(\beta^2 - N\beta - 1) - N + \frac{1}{\beta^{n+1}}\}. \end{aligned}$$

We remark that for $N = 1$, under the condition $F_1(\beta) = 0$, one sees $\hat{p}_0 = \frac{1}{(\beta-1)^2}$.

Let $p_0, e_0, f_0, e_1, f_1, \dots, e_{l-1}, f_{l-1}, q_{l-1}, r_{l-1}$ be the set of minimal projections of $\mathcal{A}_{Z_N, l}$ considered in the previous section. For a state φ on $\mathcal{A}_{Z_N}$, put

$$\hat{t} := \varphi(t) \quad \text{for } t = p_0, e_0, f_0, e_1, f_1, \dots, e_{l-1}, f_{l-1}, q_{l-1}, r_{l-1}.$$

Then φ satisfies the condition $\varphi \circ \lambda_{Z_N} = \beta\varphi$ on $\mathcal{A}_{Z_N}$ if and only if the real numbers $\hat{p}_0, \hat{e}_0, \hat{f}_0, \hat{e}_1, \hat{f}_1, \dots, \hat{e}_{l-1}, \hat{f}_{l-1}, \hat{q}_{l-1}, \hat{r}_{l-1}$ are all nonnegative and satisfy all the equations from (5.1) to (5.9) for all $l = 2, 3, \dots$. Therefore we have

Proposition 5.13. *A state φ on $\mathcal{A}_{Z_N}$ satisfies the condition $\varphi \circ \lambda_{Z_N} = \beta\varphi$ for some real number β if and only if $\beta > N$ and β is the unique solution of the equation:*

$$\beta^6 - (N+3)\beta^5 + (3N+1)\beta^4 - 2(N-1)\beta^3 - (N+2)\beta^2 + N - 1 = 0.$$

Moreover a state φ that satisfies the above condition is unique.

The topological entropy of some classes of subshifts, including irreducible topological Markov shifts, β -shifts for real number $\beta > 1$ and the subshift Z_N for $N = 1$ have appeared as the logarithm of the inverse temperature of the admitted KMS states for gauge actions on the associated C^* -algebras ([10], [14], [26]). For a subshift (Λ, σ) and a natural number k , let $\theta_k(\Lambda)$ be the cardinal number of the words $B_k(\Lambda)$ of length k appearing in Λ . The topological entropy $h_{\text{top}}(\Lambda)$ for (Λ, σ) is given by

$$h_{\text{top}}(\Lambda) = \lim_{k \rightarrow \infty} \frac{1}{k} \log \theta_k(\Lambda) \quad (\text{cf. [23]}).$$

For the subshifts Z_N , we have

Lemma 5.14. *If there exists a $\log \beta$ -KMS state on $\mathcal{O}_{Z_N}$ for gauge action for some real number $\beta > N$, we have*

$$\log \beta = \log r(\lambda_{Z_N}) = h_{\text{top}}(Z_N)$$

where $r(\lambda_{Z_N})$ denotes the spectral radius of the operator λ_{Z_N} on $\mathcal{A}_{Z_N}$.

Proof. The proof is similar to the proof of Lemma 6.7 of [26]. For the sake of completeness, we will give a proof. A word $\mu = \mu_1 \dots \mu_n$ in Σ_N appears in the subshift Z_N if and only if $S_\mu (= S_{\mu_1} \dots S_{\mu_n}) \neq 0$. Let φ be a $\log \beta$ -KMS state on $\mathcal{O}_{Z_N}$ for gauge action for some positive real number β . For $k \in \mathbb{N}$, it follows that

$$\beta^k = \varphi\left(\sum_{\mu \in B_k(Z_N)} S_\mu^* S_\mu\right) \leq \left\| \sum_{\mu \in B_k(Z_N)} S_\mu^* S_\mu \right\| = \|\lambda_{Z_N}^k(1)\| \leq \sum_{\mu \in B_k(Z_N)} \|S_\mu^* S_\mu\| = \theta_k(Z_N).$$

As $\lambda_{Z_N}^k$ is a completely positive map on the unital C^* -algebra $\mathcal{A}_{Z_N}$, we have $\|\lambda_{Z_N}^k(1)\| = \|\lambda_{Z_N}^k\|$ so that we see

$$\beta^k \leq \|\lambda_{Z_N}^k\| \leq \theta_k(Z_N). \quad (5.19)$$

On the other hand, by the inequality $\beta^k \geq \theta_k(Z_N) \min_{\mu \in Z^k} \varphi(S_\mu^* S_\mu)$, we obtain

$$\min_{\mu \in B_k(Z_N)} \varphi(S_\mu^* S_\mu)^{\frac{1}{k}} \cdot \theta_k(Z_N)^{\frac{1}{k}} \leq \beta \leq \theta_k(Z_N)^{\frac{1}{k}}.$$

Now we have $S_\mu^* S_\mu \geq P_0$ for any word $\mu \in B_*(Z_N)$. It follows that

$$\varphi(S_\mu^* S_\mu) \geq \varphi(P_0) = \frac{(\beta-2)(\beta^2 - N\beta - 1)}{N(\beta^2 - 1)} > 0 \quad \text{for } \mu \in B_*(Z_N).$$

Hence we obtain

$$\lim_{k \rightarrow \infty} \min_{\mu \in B_k(Z_N)} \varphi(S_\mu^* S_\mu)^{\frac{1}{k}} = 1$$

and $\lim_{k \rightarrow \infty} \theta_k(Z_N)^{\frac{1}{k}} = \beta$. Thus we get the desired equalities from (5.19). $\square$

Therefore we conclude

Theorem 5.15. (i) *For a positive real number β , there exists a $\log \beta$ KMS-state for gauge action on $\mathcal{O}_{Z_N}$ if and only if $\beta > N$ and β is the unique solution of the equation:*

$$\beta^6 - (N+3)\beta^5 + (3N+1)\beta^4 - 2(N-1)\beta^3 - (N+2)\beta^2 + N - 1 = 0.$$

(ii) *The above KMS-state is unique.*

(iii) $\log \beta = h_{\text{top}}(Z_N)$: *the topological entropy for the subshift Z_N .*

We finally mention an asymptotic behavior of the solution $\beta > N$ for $F_N(\beta) = 0$. Let $\beta_N > N$ be the unique solution for the equation $F_N(\beta) = 0$. We know them by numerical calculation as in the following way:

$$\begin{aligned} \beta_1 &= 2.652891650 \dots \\ \beta_2 &= 3.063607825 \dots \\ \beta_3 &= 3.670666991 \dots \\ \beta_4 &= 4.446202651 \dots \\ \beta_5 &= 5.321226229 \dots \\ \beta_6 &= 6.247124025 \dots \\ \beta_7 &= 7.199582119 \dots \\ \beta_8 &= 8.166942400 \dots \\ &\dots \end{aligned}$$

Proposition 5.16. *Let $\beta_N > N$ be the unique solution of the equation $F_N(\beta) = 0$. Then we have*

$$\lim_{N \rightarrow \infty} \frac{\beta_N}{N} = 1.$$

Proof. Put $t_N = \frac{\beta_N}{N}$. It follows that

$$\begin{aligned} 0 &= \frac{F_N(\beta_N)}{N^6} \\ &= t_N^6 - (1 + \frac{3}{N})t_N^5 + (\frac{3}{N} + \frac{1}{N})t_N^4 - 2(\frac{1}{N^2} - \frac{1}{N^3})t_N^3 - (\frac{1}{N^3} + \frac{2}{N^4})t_N^2 + \frac{1}{N^5} - \frac{1}{N^6} \end{aligned}$$

and hence

$$1 - t_N = \frac{3}{N} + (\frac{3}{N} + \frac{1}{N})\frac{1}{t_N} - 2(\frac{1}{N^2} - \frac{1}{N^3})\frac{1}{t_N^2} - (\frac{1}{N^3} + \frac{2}{N^4})\frac{1}{t_N^3} + (\frac{1}{N^5} - \frac{1}{N^6})\frac{1}{t_N^5}.$$

As $0 < \frac{1}{t_N} < 1$, we get $\lim_{N \rightarrow \infty} |t_N - 1| = 0$. $\square$

6. FLOW EQUIVALENCE CLASSES OF THE SUBSHIFTS $Z_N, N \in \mathbb{N}$

We will finally apply our discussions to a classification problem in symbolic dynamical systems under flow equivalence (cf. [3], [12], [35]). In [30], [31], we have defined the K-groups $K_i(\Lambda), i = 0, 1$ and the Bowen-Franks groups $BF^i(\Lambda), i = 0, 1$ for subshift Λ by the K-groups $K_i(\mathcal{O}_\Lambda), i = 0, 1$ and the Ext-groups $\text{Ext}^i(\mathcal{O}_\Lambda), i = 0, 1$ for the associated C^* -algebra $\mathcal{O}_\Lambda$ respectively. We have then proved that the groups $K_i(\Lambda), BF^i(\Lambda), i = 0, 1$ are invariant under not only topological conjugacy class but also flow equivalence class of subshifts. Especially there is no known computable invariant under flow equivalence of subshifts other than the groups. Since the Ext-groups $\text{Ext}^i(\mathcal{O}_\Lambda), i = 0, 1$ can be computed by the Universal Coefficient Theorem for the K-theory of C^* -algebra $\mathcal{O}_\Lambda$, we have by Theorem 4.8,

$$\begin{aligned} K_0(Z_N) &= \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}, & K_1(Z_N) &= \{0\}, \\ BF^0(Z_N) &= \mathbb{Z}/N\mathbb{Z}, & BF^1(Z_N) &= \mathbb{Z}. \end{aligned}$$

Therefore we have

Proposition 6.1. *The subshifts $Z_N, N = 1, 2, \dots$ are not flow equivalent to each other.*

Acknowledgements. The author would like to deeply thank Wolfgang Krieger for his many useful conversations.

REFERENCES

- [1] M. P. BÉAL, *Codage Symbolique*, Masson, Paris (1993).
- [2] F. BLANCHARD AND G. HANSEL, *Systems codés*, Theor. Computer Sci. **44** (1986) pp. 17–49.
- [3] R. BOWEN AND J. FRANKS, *Homology for zero-dimensional nonwandering sets*, Ann. Math. **106** (1977) pp. 73–92.
- [4] T. M. CARLESEN AND K. MATSUMOTO, *Some remarks on the C^* -algebras associated with subshifts*, Math. Scand. **95**(2004), pp. 145–160.
- [5] N. CHOMSKY AND M. P. SCHÜTZENBERGER, *The algebraic theory of context-free languages*, Computer programming and formal systems, North-Holland (1963), pp. 118–161.
- [6] J. CUNTZ, *Simple C^* -algebras generated by isometries*, Comm. Math. Phys. **57**(1977) pp. 173–185.
- [7] J. CUNTZ, *A class of C^* -algebras and topological Markov chains II: reducible chains and the Ext-functor for C^* -algebras*, Inventiones Math. **63**(1980) pp. 25–40.
- [8] J. CUNTZ, *K-theory for certain C^* -algebras*, Ann. Math. **117**(1981) pp. 181–197.
- [9] J. CUNTZ AND W. KRIEGER, *A class of C^* -algebras and topological Markov chains*, Invent. Math. **56**(1980), pp. 251–268.
- [10] M. ENOMOTO, M. FUJII AND Y. WATATANI, *KMS states for gauge action on $\mathcal{O}_A$* , Math. Japon **29** (1984) pp. 607–619.
- [11] R. FISCHER, *Sofic systems and graphs*, Monats. für Math. **80** (1975) pp. 179–186.
- [12] J. FRANKS, *Flow equivalence of subshifts of finite type*, Ergodic Theory Dynam. Systems **4** (1984) pp. 53–66.
- [13] J. E. HOPCROFT AND J. D. ULLMAN, *Introduction to Automata Theory, Languages, and Computation*, Addison-Wesley, Reading (2001).
- [14] Y. KATAYAMA, K. MATSUMOTO AND Y. WATATANI, *Simple C^* -algebras arising from β -expansion of real numbers*, Ergodic Theory Dynam. Systems **18** (1998) pp. 937–962.
- [15] E. KIRCHBERG, *The classification of purely infinite C^* -algebras using Kasparov's theory*, preprint,(1994).
- [16] B. P. KITCHENS, *Symbolic dynamics*, Springer-Verlag, Berlin, Heidelberg and New York (1998).
- [17] W. KRIEGER, *On sofic systems I*, Israel J. Math. **48**(1984), pp. 305–330.
- [18] W. KRIEGER, *On sofic systems II*, Israel J. Math. **60**(1987), pp. 167–176.

- [19] W. KRIEGER, *On subshifts and topological Markov chains*, Numbers, information and complexity (Bielefeld 1998), Kluwer Acad. Publ. Boston MA (2000) pp. 453–472
- [20] W. KRIEGER AND K. MATSUMOTO, *Shannon graphs, subshifts and lambda-graph systems*, J. Math. Soc. Japan **54**(2002), pp. 877–900.
- [21] W. KRIEGER AND K. MATSUMOTO, *Subshifts from certain one-counter codes*, preprint.
- [22] M. LACA AND S. NESHVEYEV, *KMS-states of quasi-free dynamics on Pimsner algebras*, J. Funct. Anal. **211** (2004) pp. 457–482.
- [23] D. LIND AND B. MARCUS, *An introduction to symbolic dynamics and coding*, Cambridge University Press, Cambridge (1995).
- [24] K. MATSUMOTO, *On C^* -algebras associated with subshifts*, Internat. J. Math. **8**(1997), pp. 357–374.
- [25] K. MATSUMOTO, *K-theory for C^* -algebras associated with subshifts*, Math. Scand. **82**(1998), pp. 237–255.
- [26] K. MATSUMOTO, *A simple C^* -algebra arising from certain subshift*, J. Operator Theory **42** (1999), pp. 351–370.
- [27] K. MATSUMOTO, *Dimension groups for subshifts and simplicity of the associated C^* -algebras*, J. Math. Soc. Japan **51** (1999), pp. 679–698.
- [28] K. MATSUMOTO, *Presentations of subshifts and their topological conjugacy invariants*, Documenta Math. **4** (1999), pp. 285–340.
- [29] K. MATSUMOTO, *Relations among generators of C^* -algebras associated with subshifts*, Internat. J. Math. **10**(1999), pp. 385–405.
- [30] K. MATSUMOTO, *Bowen-Franks groups for subshifts and Ext-groups for C^* -algebras*, K-Theory **23** (2001), pp. 67–104.
- [31] K. MATSUMOTO, *Bowen-Franks groups as an invariant for flow equivalence of subshifts*, Ergodic Theory Dynam. Systems **21** (2001), pp. 1831–1842.
- [32] K. MATSUMOTO, *C^* -algebras associated with presentations of subshifts*, Documenta Math. **7** (2002), pp. 1–30.
- [33] K. MATSUMOTO, *Construction and pure infiniteness of C^* -algebra associated with lambda-graph systems*, Math. Scand. **97** (2005), pp. 73–89.
- [34] K. MATSUMOTO, Y. WATATANI AND M. YOSHIDA, *KMS-states for gauge actions on C^* -algebras associated with subshifts*, Math. Z. **228** (1998) pp. 489–509.
- [35] W. PARRY AND D. SULLIVAN, *A topological invariant for flows on one-dimensional spaces*, Topology **14** (1975) pp. 297–299.
- [36] C. PINZARI, Y. WATATANI AND K. YOSHIDA, *KMS-states, entropy and the variational principle in full C^* -dynamical systems*, Comm. Math. Phys. **213** (2000) pp. 231–379.
- [37] N. C. PHILLIPS, *A classification theorem for nuclear purely infinite simple C^* -algebras*, Doc. Math. **5** (2000) pp. 49–114.
- [38] M. RØDOM, *Classification of Cuntz-Krieger algebras*, K-theory **9**(1995), pp. 31–58.
- [39] M. RØDOM, *Classification of purely infinite simple C^* -algebras I*, J. Func. Anal. **131**(1995), pp. 415–458.
- [40] B. WEISS, *Subshifts of finite type and sofic systems*, Monats. Math. **77**, (1973), pp. 462–474.

DEPARTMENT OF MATHEMATICAL SCIENCES, YOKOHAMA CITY UNIVERSITY, 22-2 SETO, KANAZAWA-KU, YOKOHAMA, 236-0027 JAPAN
E-mail address: kengo@yokohama-cu.ac.jp